



PERSIAN GULF METHANOLIC EXTRACT OF *NERITA LONGI* AS EFFICIENT ANTIOXIDANT AGENT

Mohammad Reza Shushizadeh^{1,2*}, Khosnavaz Farzan¹, Nabavi Mohammad Bagher^{2,3}, Rahbar Nadere¹, Namjooyan Forough¹

1. *Department of Medicinal Chemistry, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, IR Iran.*
2. *Marine Pharmaceutical Science Research Center, Ahvaz Jundishapur University of Medical Sciences, IR Iran.*
3. *Department of Marine Biology, Marine Science and Technology University of Khoramshahr, IR Iran.*

ARTICLE INFO

Received:

03th Jun 2017

Accepted:

29th Nov 2017

Available online:

14th Dec 2017

Keywords:

ABSTRACT

Background: Marine organisms are the important key for the development and conception of new drugs with medicine applications. Marine invertebrates produce a high amount of natural products as adaptation to their environmental and as result of their life style. Marine natural compounds with powerful bioactivities were already extracted from marine invertebrates including gastropods, tunicates, sponges, soft corals, sea hares, sea cucumbers and bryozoans. Some of those compounds, have pharmaceutical activity such as antitumor, antimicrobial, anti-inflammatory and antioxidant activities which can be used in clinical and preclinical assays.

Objectives: The purpose of this research was to examine the activity of methanolic extract from Persian Gulf gastropod, *Nerita longi* as efficient antioxidant.

Materials and methods: Persian Gulf gastropod, *Nerita longi* was washed and macerated with ethyl acetate and methanol for three days. The resulting extracts were dry with freeze dryer and the final residue were consider for antioxidant tests by DPPH, FRAP and FOLIN methods.

Results: In DPPH method, the resultant IC₅₀ for methanol and ethyl acetate were 8.07 and 10.13 mg/ml, respectively. In FRAP method, the resultant EC₁ for methanol and ethyl acetate were 3.687, 5.35 mg/ml respectively. In FOLIN method, the highest Absorbance in methanolic and ethyl acetate extracts are given 1.626 and 1.592.

Conclusions: Muscle tissue of Persian Gulf gastropod, *Nerita Longi*, is an important source of natural proteins and peptides. So high antioxidant activity was given from FRAP, DPPH and FOLIN methods of methanolic extract can be related to the ability of methanol to extract corresponding peptides and proteins.

Copyright © 2013 - All Rights Reserved - Pharmacophore

To Cite This Article: Mohammad Reza Shushizadeh, Khosnavaz Farzan, Nabavi Mohammad Bagher, Rahbar Nadere, Namjooyan Forough, (2017), "Persian Gulf methanolic extract of *Nerita longi* as efficient antioxidant agent", *Pharmacophore*, **8(6S)**, e-1173266.

Introduction

MATERIALS AND METHODS

Chemicals

2,2-diphenyl-2-picrylhydrazyl (DPPH), 2,4,6-tripyridyl-s-triazine (TPTZ) and ferulic acid were purchased from Sigma Chemical Co. (St. Louis, MO, USA); Folin–Ciocalteu reagent, ferric chloride (FeCl₃ · 6H₂O), and methanol and ethyl acetate were from Merck (Darmstadt, Germany)

Sample collection and extraction

Live specimens of *Nerita Longii* gastropod was collected from the Bushehr coast of Persian Gulf of Iran and directly brought to the laboratory and identified by scientists from Marine Science and Technology of Khoramshahr, Khuzestan, Iran. The whole body muscle was taken out from the shell, used for extraction by methanol and stored for 4 days.

Preparation of extracts

The shells were separated in lab and the whole muscle tissue weighing 80 g was macerated with ethyl acetate and methanol for three days. The supernatant solvent was concentrated using vacuum evaporator (35-55°C) under reduced pressure and the resultant extract was dry with freeze dryer and kept in clean glass vials at -80 °C until use.

DPPH radical scavenging assay

The free radical scavenging activity of methanol and ethyl acetate extract of *N. Longi* was assessed using DPPH (1,1-diphenyl-2-picryl hydrazyl) radical(1). Due to the poor solubility of methanol and ethyl acetate extract in aqueous medium it was dissolved in 1 % DMSO and later on made up to desired concentrations with distilled water. The extract in different concentrations were added to 0.1 ml of 1 M Tris–HCl (pH 7.9), and then mixed with 0.6 mL of DPPH (100 μM) in methanol for 20 min at room temperature under protection from light. Absorbance of the mixture was read on a UV-VIS spectrophotometer (Hitachi, U-2001) at 517nm against the blank. Butylated Hydroxyl Toluene (BHT) was used as the standard antioxidant and the percentage inhibition of test and the standard were calculated as

$$\% \text{ Inhibition} = ((A_{\text{Control}} - A_{\text{Test}}) / A_{\text{Control}}) \times 100$$

Where A_{Control} is the absorbance of the control (without extract) and A_{Test} is the absorbance of the sample of extract and standard.

Ferric Ion Reducing Antioxidant Power Assay


The FRAP assay also takes advantage of electron-transfer reactions. Herein a ferric salt, Fe(III)(TPTZ)₂Cl₃ (TPTZ) 2,4,6-tripyridyls-triazine), is used as an oxidant (Benzie et al, 1996). The FRAP assay involves the following procedures: The oxidant in the FRAP assay is prepared by mixing TPTZ (2.5 mL, 10 mM in 40 mM HCl), 25 mL of acetate buffer, and 2.5 mL of FeCl₃·6H₂O (20 mM). The conglomerate is referred to as “FRAP reagent”. The final solution has Fe(III) of 1.67 mM and TPTZ of 0.83 mM. Therefore, the TPTZ is deficient as the ideal reaction stoichiometry between Fe(III) and TPTZ is 1 to 2. The oxidant is not just Fe(III)(TPTZ)₂, it also contains other Fe(III) species which can lead to potential problems as many metal chelators in food extract could bind Fe(III) and form complexes that are also capable of reacting with antioxidants. To measure FRAP value, 300 μL of freshly prepared FRAP reagent is warmed to 37 °C and a reagent blank reading is taken at 593 nm; then 10 μL of sample and 30 μL of water are added.

Absorbance readings are taken after 0.5 s and every 15 s until 4 min. The change of absorbance (ΔA) A_{4min} - A_{0min}) is calculated and related to ΔA of an Fe(II) standard solution. ΔA is linearly proportional to the concentration of antioxidant. One FRAP unit is arbitrarily defined as the reduction of 1 mol of Fe(III) to Fe(II). The FRAP values for ascorbic acid, R-tocopherol, and uric acid are identical (2.0). The FRAP value of bilirubin is 1-fold higher than that of ascorbic acid. These results suggest that 1 mol of vitamin C can reduce 2 mol of Fe(III) and that 1 mol of bilirubin can reduce 4 mol of Fe(III). This is in conflict with the fact that both vitamin C and bilirubin are two-electron reductants. It is known that when bilirubin is oxidized, it is transformed to beliverdin (by losing two hydrogen atoms, not just electrons), which happens to have an absorption at 593 nm with coefficient (593) comparable with that of Fe(II)(TPTZ)₂. Pulido and co-workers measured the FRAP values of several polyphenols in water and methanol. However, the absorption (A₅₉₃) does not stop at 4 min; instead, it slowly increased even after several hours. Polyphenols with such behaviors include caffeic acid, tannic acid, ferulic acid, ascorbic acid, and quercetin. The FRAP values of these compounds cannot be obtained accurately if 4 min reaction time was followed.

Total Phenols Assay by Folin-Ciocalteu Reagent.

FCR was initially intended for the analysis of proteins taking advantage of the reagent's activity toward protein tyrosine (containing a phenol group) residue (2). Many years later, Singleton and co-workers extended this assay to the analysis of total phenols in wine; since then the assay has found many applications. The FCR-based assay gained popularity and is commonly known as the total phenols (or phenolic) assay. The FCR actually measures a sample's reducing capacity, but this is not reflected in the name "total phenolic assay". Numerous publications applied the total phenols assay by FCR and an ET-based antioxidant capacity assay (e.g., FRAP, TEAC, etc.) and often found excellent linear correlations between the "total phenolic profiles" and "the antioxidant activity". This is not surprising if one considers the similarity of chemistry between the two assays. One of the assays may just be redundant. A recent report of using polyphenol oxidase for assaying total phenols in tea may be more specific to phenolic compounds.

The FCR is typically made by first boiling (for 10 h) the mixture of sodium tungstate ($\text{Na}_2\text{WO}_4 \cdot 2\text{H}_2\text{O}$, 100 g), sodium molybdate ($\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$, 25 g), concentrated hydrochloric acid (100 mL), 85% phosphoric acid (50 mL), and water (700 mL). After boiling, lithium sulfate ($\text{Li}_2\text{SO}_4 \cdot 4\text{H}_2\text{O}$, 150 g) is added to the mixture to give an intense yellow solutions the FC reagent. Contamination of reductants leads to a green color, and the addition of oxidants such as bromine can restore the desired yellow color. The exact chemical nature of the FC reagent is not known, but it is believed to contain heteropolyphosphotunstates- molybdates. Sequences of reversible one- or two-electron reduction reactions lead to blue species, possibly $(\text{PMoW}_{11}\text{O}_{40})^{4-}$. In essence, it is believed that the molybdenum is easier to be reduced in the complex and electron-transfer reaction occurs between reductants and Mo(VI):



Obviously, the FC reagent is nonspecific to phenolic compounds as it can be reduced by many non-phenolic compounds [e.g., vitamin C, Cu (I), etc.]. Phenolic compounds react with FCR only under basic conditions (adjusted by a sodium carbonate solution to $\text{pH}=10$). Dissociation of a phenolic proton leads to a phenolate anion, which is capable of reducing FCR. This supports the notion that the reaction occurs through electron transfer mechanism. The blue compounds formed between phenolate and FCR are independent of the structure of phenolic compounds, therefore ruling out the possibility of coordination complexes formed between the metal center and the phenolic compounds.

Despite the undefined chemical nature of FCR, the total phenols assay by FCR is convenient, simple, and reproducible.

As a result, a large body of data has been accumulated, and it has become a routine assay in studying phenolic antioxidants.

Statistical analysis

Statistical analysis was done using SPSS version 14. IC₅₀ values respectively for DPPH were found out by Linear Regression Probit analysis.

RESULTS AND DISCUSSION

DPPH radical scavenging activity

DPPH radical scavenging property of *N. Longi* is shown in Figure 1. The methanolic extract showed a dose dependent pattern in DPPH radical scavenging in the range of concentrations tested (0.312 – 20 mg/mL). The IC₅₀ value of the methanolic extract was found to be 8.07 mg/mL. The higher concentration of methanol extract showed a percentage inhibition of 82.86 %.




Figure 1: Effect of methanol extract of *N. Longi* on DPPH radical scavenging activity.

As showed in Figure 2, the EtOAc extract have a dose dependent pattern in DPPH radical scavenging in the range of their concentrations (0.312 – 20 mg/mL). The IC₅₀ value of the EtOAc extract was found to be 10.13 mg/mL. The higher concentration of EtOAc extract showed a percentage inhibition of 76.04 %.




Figure 2: Effect of EtOAc extract of *N. Longi* on DPPH radical scavenging activity.

As showed in Figure 3, the resultant IC₅₀ for methanol and ethyl acetate were 8.07 and 10.13 mg/ml, respectively.




Figure 3. Effect of IC₅₀ on Methanol and EtOAc extract of *N. Longi* on DPPH radical scavenging activity

Ferric reducing activity based on FRAP assay

The ethanolic extract of Sulawesian beans exhibited the highest antioxidant potential among the extracts (Figure 6), based on the FRAP assay. FRAP assay measures the reducing potential of an antioxidant reacting with a ferric tripyridyltriazine (Fe^{3+} -TPTZ) complex and producing a colored ferrous tripyridyltriazine (Fe^{2+} -TPTZ) (3). Generally, the reducing properties are associated with the presence of compounds, which exert their action by breaking the free radical chain through donating a hydrogen atom (4). According to Benzie and Strain (3), the reduction of Fe^{3+} -TPTZ complex to blue colored of Fe^{2+} -TPTZ occurs at low pH.

Sulawesian and Malaysian beans are well known to have a low cotyledon pH, while Ghanaian has medium pH, and Ivory Coast has high pH (5). The highest antioxidant potential of Malaysian beans could be due to the highly acidic (low pH) nature of the bean cotyledon, which many influence the pH of the assay medium.

As depicted in the Figure 4, the methanolic extract showed a dose dependent pattern in FRAP scavenging in the range of their concentrations (0.312–10 mg/ml). The EC1 value of the methanolic extract was found to be 3.687 mg/mL. The higher Absorbance of methanolic extract showed at 1.78.




Figure 4. Effect of methanolic extract on Absorbance of *N. Longi* in FRAP Value

As depicted in Figure 5, the EtOAc extract showed a dose dependent pattern in DPPH radical scavenging in the range of concentrations tested (0.312 – 20 mg/mL). The EC1 value of the EtOAc extract was found to be 5.35 mg/mL. The higher concentration of EtOAc extract showed the absorbance at 1.822




Figure 5. Effect of EtOAc extract on Absorbance of *N. Longi* in FRAP Value

As shown in Figure 6, the resultant EC₁ for methanol and ethyl acetate were 3.687, 5.35 mg/mL respectively.




Figure 6: The comparison of EC₁ in methanolic and EtOAc extract of FRAP Value

Total phenolic content

Several studies showed a correlation between antioxidant activity and phenolic content (6; 7; 8). As shown in Figure 7, methanolic extract beans had the highest phenolic content. The highest Absorbance in methanolic extract is 1.626.




Figure 7: Effect of methanolic extract on Absorbance of N. Longi in FOLIN

As shown in Figure 8 the highest Absorbance in EtOAc is 1.592.




Figure 8: Effect of EtOAc extract on Absorbance of *N. Longi* in FOLIN

Conclusion

Muscle tissue of Persian Gulf gastropod, *Nerita Longi*, is an important source of natural proteins and peptides. So high antioxidant activity was given from FRAP, DPPH and FOLIN methods of methanolic extract can be related to the ability of methanol to extract corresponding peptides and proteins.

Acknowledgements:

The authors wish to thank College of Pharmacy and Marine Pharmaceutical Research Center of Ahvaz Jundishapur University of Medical Science for financial support of this work.

References:

1. Jayaprakasha G.K., Jaganmohan Rao L., Sakariah K.K. (2004). Antioxidant activities of flavidin in different in vitro model systems, *Bioorg. Med. Chem.* 12, 5141-5146.
2. Woraratphoku J, Intarapichet KO, Indrapichate A. (2007) Phenolic compounds and antioxidative properties of selected Wines from the northeast of Thailand. *Food Chemistry*, 104, 1485-1490.
3. Benzie IFF, Strain JJ, The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": The FRAP assay, *Analytical Biochemistry* 1996.; 239 (1), 70-76.
4. Gordon, M. H. (1990). The mechanism of antioxidant action in-vitro. In B. J. F. Hudson (Ed.), *Food antioxidants* (pp. 1-18). London: Elsevier Applied Science.
5. Misnawi Jinap, S., Nazamid, S., Jamilah, B. (2002). Activation of remaining key enzymes in dried under-fermented cocoa beans and its effect on aroma precursor formation. *Food Chemistry*, 78, 407-417.
6. Nagai, T., Reiji, I., Hachiro, I., Nobutaka, S. (2003). Preparation and antioxidant properties of water extract of propolis. *Food Chemistry*, 80, 29-33.
7. Velioglu, Y. S., Mazza, G., Gao, L., Oomah, B. D. (1998). Antioxidant activity and total phenolics in selected fruits, vegetables and grain products. *Journal of Agricultural and Food Chemistry*, 46, 4113-4117.
8. Yang, J. H., Lin, H. C., Mau, J. L. (2002). Antioxidant properties of several commercial mushrooms. *Food Chemistry*, 77, 229-235.